

Local Best Rational Approximations to Continuous Functions and the Rays They Emanate

DANIEL WULBERT

*Mathematics Department, University of California at San Diego,
La Jolla, California 92093, U.S.A.*

Communicated by E. W. Cheney

Received February 15, 1985

Let f be a continuous real function defined on $[0, 1]$. A real rational function $r_0 \in R_n^m(\mathbb{C})$ is a local best approximation to $cf + (1 - c)r_0$ for each $c > 0$ if and only if r_0 is a global best approximation to f from $\text{Re } R_n^m(\mathbb{C})$. © 1988 Academic Press, Inc.

I. INTRODUCTION

Suppose that a real rational function r_0 is a local best approximation to a continuous real function f from the real rational functions R_n^m . It is known then that r_0 is a global best approximation to f , and that it also is a best approximation to each function on the ray $\{f_c = cf + (1 - c)r_0: c \geq 0\}$. However, if r_0 is the best approximation from $R_n^m(\mathbb{C})$ —the complex valued rationals defined on the unit interval—it is not necessarily a best approximation to each f_c . Moreover it is not known, in the complex setting, if r_0 being a local best approximation implies that it is a global best approximation.

We show here that r_0 being a local best approximation to all f_c from $R_n^m(\mathbb{C})$ is a very strong condition, equivalent to r_0 being a global best approximation from $\text{Re } R_n^m(\mathbb{C})$.

Notation. The real polynomials of degree less than or equal to k which are defined on $[0, 1]$ are denoted by \mathcal{P}_k . The corresponding complex polynomials are written $\mathcal{P}_k(\mathbb{C})$. The degree of a polynomial p is ∂p .

$$\mathcal{P}_k^+ = \{p \in \mathcal{P}_k: p(x) \neq 0 \text{ for } 0 \leq x \leq 1\}, \quad (1.1)$$

and

$$\mathcal{R}_n^m = \{p/q: p \in \mathcal{P}_m, q \in \mathcal{P}_n^+\}. \quad (1.2)$$

Analogous statements define $\mathcal{P}_k^+(\mathbb{C})$ and $\mathcal{R}_n^m(\mathbb{C})$.

For a function g and a set $K \subseteq [0, 1]$,

$$\|g\|_K = \sup\{|g(k)|: k \in K\}, \quad (1.3)$$

and

$$\|g\| = \|g\|_{[0,1]}. \quad (1.4)$$

We use

$$\text{crit } g = \{x \in [0, 1]: |g(x)| = \|g\|\}, \quad (1.5)$$

and

$$\text{sgn } g = \begin{cases} \frac{g(x)}{|g(x)|}, & x \neq 0 \\ 0, & x = 0. \end{cases} \quad (1.6)$$

As usual $\text{Re } g$ and $\text{Im } g$ represent the real and imaginary parts of g .

For a set, A , of functions on $[0, 1]$,

$$\begin{aligned} \text{Re } A &= \{\text{Re } g: g \in A\} \\ \text{Im } A &= \{\text{Im } g: g \in A\}. \end{aligned} \quad (1.7)$$

A function f is said to have $g \in A$ as a *best approximation* from A if

$$\|f - g\| = \inf\{\|f - a\|: a \in A\}. \quad (1.8)$$

If there is a neighborhood U of g such that g is a best approximation to f from $A \cap U$, then g is a *local best approximation* to f .

Reserved Notation. We will reserve the following notation throughout the paper, $p_0 \in \mathcal{P}_m$, $q_0 \in \mathcal{P}_n$. We assume that p_0 and q_0 have no common factors.

$$r_0 = p_0/q_0, \quad (1.9)$$

and

$$d = \max\{m - \partial p_0, n - \partial q_0\}. \quad (1.10)$$

Let f be a continuous real function on $[0, 1]$; we write, for c real,

$$f_c = cf + (1 - c)r_0, \quad (1.11)$$

$$e_c = f_c - r_0 \quad \text{and} \quad e = e_1. \quad (1.12)$$

II. ESTIMATES FOR $\|e_c\|$

The proof of the main theorem uses numerous computations. This section collects results which conclude that a function g has the property that $\|e_c - g\| < \|e_c\|$.

LEMMA 2.1. *If*

$$\|e - g\|_{\text{crit } e} < \|e\|_{\text{crit } e},$$

then for large c ,

$$\|e_c - g\| < \|e_c\|.$$

Proof. There is a neighborhood U of $\text{crit } e$ for which

$$\|e - g\|_U < \|e\|. \quad (2.1)$$

Hence

$$\begin{aligned} \|e_c - g\|_U &\leq \|(c-1)(f - r_0)\|_U + \|f - r_0 - g\|_U \\ &\leq (c-1) \|e\|_U + \|e\| \\ &\leq c \|e\| \\ &= \|e_c\|. \end{aligned} \quad (2.2)$$

For points not in U we have an $\varepsilon > 0$ such that

$$\|e\|_{[0,1] - U} \leq \|e\| - \varepsilon. \quad (2.3)$$

So in this case,

$$\begin{aligned} \|e_c - g\|_{[0,1] - U} &\leq \|e_c\|_{[0,1] - U} + \|g\| \\ &\leq c[\|e\| - \varepsilon] + \|g\|. \end{aligned} \quad (2.4)$$

So if

$$\begin{aligned} c &> \|g\|/\varepsilon, \\ \|e_c - g\|_{[0,1] - U} &\leq c \|e\| = \|e_c\|. \end{aligned} \quad (2.5)$$

Combining (2.3) and (2.5) proves the lemma. ■

LEMMA 2.2. *If*

$$\|e - \text{Re } g\| < \|e\|,$$

then for large c

$$\|e_c - g\| < \|e_c\|.$$

Proof. From Lemma 2.1 we need only show that

$$\|e - g\|_{\text{crit } e} < \|e\|. \quad (2.6)$$

For x in $\text{crit } e = \text{crit } e_c$,

$$|(e_c - g)(x)|^2 < \|e_c\|^2, \quad (2.7)$$

if and only if

$$\|e_c\|^2 - 2ce(x) \operatorname{Re} g(x) + [\operatorname{Re} g(x)]^2 + [\operatorname{Im} g(x)]^2 \leq \|e_c\|^2, \quad (2.8)$$

if and only if

$$|(e - g)(x)|^2 - (c - 1)e(x) \operatorname{Re} g(x) + [\operatorname{Im} g(x)]^2 \leq \|e\|^2, \quad (2.9)$$

if and only if

$$[\operatorname{Im} g(x)]^2 \leq (c - 1)e(x). \quad (2.10)$$

From the hypothesis, $\operatorname{Re} g(x)$ must be a nonzero number of the same sign as $e(x)$. Therefore the right side of the inequality can be made arbitrarily large with c . ■

LEMMA 2.3. *If g is a real valued function such that $\operatorname{sgn}[g(x)] = \operatorname{sgn}[e(x)]$ for all x in $\text{crit } e$, then for all sufficiently large c ,*

$$\|e_c - g\| < \|e_c\|.$$

Proof. If $\|g\|_{\text{crit } e} < \|e_c\|$, then $\|e_c - g\|_{\text{crit } e} < \|e_c\|$. Hence the lemma follows from Lemma 2.1. ■

LEMMA 2.4. $x \in \text{crit } e$, then

$$\|e\|^2 - |e(x) - g(x)|^2 = 2e(x) \operatorname{Re} g(x) - |g(x)|^2.$$

Proof. This is acquired by just expanding $|e(x) - g(x)|^2$. ■

III. CLASSES OF POLYNOMIALS

LEMMA 3.1.

$$P_0 \mathcal{P}_n + q_0 \mathcal{P}_m = \mathcal{P}_{m+n-d}.$$

Proof. A proof for this known result can be built using the degrees of the polynomials and the dimensions of the linear spaces. (For example, see [1].) ■

Notation. For a complex j times differential function f put

$$Z(f) = \{\omega \in \mathbb{C}: f(\omega) = 0\},$$

$$Z_2(f) = \{\omega \in Z(f): f'(\omega) = 0\},$$

and

$$Z_j(f) = \{\omega \in Z_{j-1}(f): f^{(j)}(\omega) = 0\}.$$

LEMMA 3.2.

$\{\gamma^2 + q_0 \mathcal{P}_n: \gamma \in \mathcal{P}_n\} \supset \{t \in \mathcal{P}_{2n}: (1) Z(t) \cap Z_2(q_0) = \emptyset, (2)$
If $\partial t > n + \partial q_0$, then t is even and t has a positive leading
coefficient, and (3) $t \geq 0$ on $Z(q_0) \cap R$ }. $\}$

Proof. We wish to find a $\gamma \in \mathcal{P}_n$ so that γ^2 agrees with t on the zeros of q_0 , and which has coefficients that agree with those of t for powers of x greater than $n + \partial q_0$. For such a γ , $t - \gamma^2$ is a polynomial of degree $n + \partial q_0$ which has q_0 as a factor and the lemma will be proven.

Let $H \in \mathcal{P}_{\partial q_0}$ be chosen so that H^2 agrees with t on the zeros of q_0 —including multiple zeros. For example, on a double zero of q_0 , the derivative of H^2 agrees with that of t . This Hermite-type interpolation is possible since H is not zero on a multiple zero of q_0 . (We use conditions (1) and (3) hypothesised for t in defining H .)

We now wish to find $S \in \mathcal{P}_{n - \partial q_0}$ so that

$$(Sq_0 + H)^2 = \gamma^2 \tag{3.1}$$

has coefficients of $X^{n + \partial q_0 + 1}, X^{n + \partial q_0 + 2}, \dots, X^{2n}$ that agree with those of t . If $It \leq n + \partial q_0$ this is satisfied with S equal zero. Hence we will assume that $\partial t = 2k > n + \partial q_0$. We proceed by examining the coefficients in the expansion of $(Sq_0 + H)^2$ (the coefficients of q_0 and H are already fixed: those for S are to be determined). For j larger than $k - \partial q_0$ put the coefficient of x^j for S equal to zero.

The leading coefficient of $(Sq_0 + H)^2$ is the product of the squares of the leading coefficients of S and q_0 . Since the leading coefficient of t is positive, the coefficient of $x^{k - \partial q_0}$ for S is determined. If $0 < j < k - \partial q_0$, the coefficient of $x^{2k - j}$ in the expansion of $(Sq_0 + H)^2$ can be written as the sum of two terms. One consists of twice the product of the lead coefficients of S , q_0 , and s_i the coefficient of $x^{\partial q_0 - i}$ in S . The other is an expression composed of coefficients of q_0 and already determined coefficients of S . Hence s_j can be chosen so that the coefficients of $x^{2k - j}$ in t and $(Sq_0 + H)^2$ are equal. ■

COROLLARY 3.3.

$$\{\gamma^2 + q_0 \mathcal{P}_n: \gamma \in \mathcal{P}_n\} \supseteq \{q^2 + s^2: q, s \in \mathcal{P}_n, Z(q^2 + s^2) \cap Z(q_0) = \emptyset\}$$

LEMMA 3.4.

$$\begin{aligned} & \{\omega(p_0\beta - q_0\alpha)q_0 - p_0\gamma^2 + q_0\gamma\delta: \omega \in \mathcal{P}_d, \alpha, \delta \in \mathcal{P}_m, \beta, \gamma \in \mathcal{P}_n\} \\ & \supseteq \{\mathcal{P}_{m+n}q_0 - p_0(q^2 + s^2): q, s \in \mathcal{P}_n, \text{ and } Z(y^2 + s^2) \cap Z(q_0) = \emptyset\}. \end{aligned}$$

Proof. From Corollary 3.3, each member of this set can be written in the form $q_0v - p_0\gamma^2$ for some $v \in \mathcal{P}_{m+n}$. Choose δ so that $v - \gamma\delta$ has a real zero. Then there are $\omega \in \mathcal{P}_d$ and $u \in \mathcal{P}_{m+n-d}$ such that

$$\omega u = v - \gamma\delta. \quad (3.2)$$

Now choose α and β so that

$$p_0\beta - q_0\alpha = u. \quad (3.3)$$

We have

$$\begin{aligned} \omega(p_0\beta - q_0\alpha)q_0 - p_0\gamma^2 + q_0\gamma\delta &= \omega u q_0 - p_0\gamma^2 + q_0\gamma\delta \\ &= v q_0 - \gamma\delta q_0 - p_0\gamma^2 + q_0\gamma\delta \\ &= v q_0 - p_0\gamma^2. \quad \blacksquare \end{aligned} \quad (3.4)$$

IV. NOTATIONAL CONVENTION

Let $\omega \in d$, $\alpha, \delta \in \mathcal{P}_m$, and $\beta, \gamma \in \mathcal{P}_n$. For the remainder of the paper we will write, for λ real,

$$r_\lambda = \frac{\omega p_0 + \lambda^2 \alpha + i\lambda \delta}{\omega q_0 + \lambda^2 \beta + i\lambda \gamma} \quad (4.1)$$

$$r = r_1 \quad (4.2)$$

$$L_\lambda = \frac{\lambda^2[\beta p_0 - \alpha q_0] \omega q_0 - \gamma^2 p_0 + \gamma \delta q_0 + i\lambda[\gamma p_0 - \delta q_0] \omega q_0}{q_0^3 \omega^2} \quad (4.3)$$

and

$$L = L_1. \quad (4.4)$$

V. LOCAL BEST APPROXIMATIONS

The next two lemmas record the result of straightforward computation from definitions.

LEMMA 5.1.

$$r_0 - r_\lambda = \lambda^2 \left\{ \frac{[\beta p_0 - \alpha q_0][\omega q_0 + \lambda^2 \beta] - [\gamma^2 p_0 - \delta \gamma q_0]}{q_0[(\omega q_0 + \lambda^2 \beta)^2 + (\lambda \gamma)^2]} \right\} \\ + i\lambda \left\{ \frac{(\gamma p_0 - \delta q_0)(\omega q_0 + \lambda^2 \beta) - \lambda^2 \gamma [\beta p_0 - \alpha q_0]}{q_0[(\omega q_0 + \lambda^2 \beta)^2 + (\lambda \gamma)^2]} \right\}.$$

LEMMA 5.2.

$$\lim_{\lambda \rightarrow 0} \left\| \frac{r_0 - r_\lambda - L_\lambda}{\lambda^2} \right\| = 0.$$

LEMMA 5.3. *If*

$$\|e - L\|_{\text{crit } e} < \|e\|,$$

then for all sufficiently small λ

$$\|e - r_\lambda\| < \|e\|.$$

Proof. From Lemma 2.4 there is an $\varepsilon > 0$ such that on $\text{crit } e$

$$2e[\text{Re } L] > |L|^2 + \varepsilon. \quad (5.1)$$

This inequality must also hold on some neighborhood U of $\text{crit } e$. It is also true that on U for $0 < \lambda \leq 1$,

$$2e\lambda^2 \text{Re } L > \lambda^4[\text{Re } L]^2 + \lambda^2[\text{Im } L]^2 + \lambda^2\varepsilon. \quad (5.2)$$

It is always true (because e is real) that

$$|e - L_\lambda|^2 \leq \|e\|^2 - 2e \text{Re } L_\lambda + |L_\lambda|^2. \quad (5.3)$$

Since

$$\lambda^2 \text{Re } L = \text{Re } L_\lambda \quad \text{and} \quad \lambda \text{Im } L = \text{Im } L_\lambda, \quad (5.4)$$

line (5.2) shows that on U ,

$$|e - L_\lambda|^2 \leq \|e\|^2 - \lambda^2\varepsilon \\ \leq \left[\|e\| - \lambda^2 \frac{\varepsilon}{2\|e\|} \right]^2. \quad (5.5)$$

From Lemma 5.2 we conclude that for small positive λ

$$\|f - r_\lambda\|_U < \|e\|. \quad (5.6)$$

Since r_λ converges uniformly to r_0 , and since there is a positive μ for which

$$\|e\|_{[0,1]} - \mu < \|e\|, \quad (5.7)$$

we obtain that for all sufficiently small λ ,

$$\|f - r_\lambda\|_{[0,1]} - \mu < \|e\|. \quad (5.8)$$

This combines with line (5.6) to prove the lemma. ■

THEOREM. *If r_0 is a local best approximation to f_c for all $c > 0$ then r_0 is a global best approximation to f from $\text{Re } R_n^m(\mathbb{C})$.*

Proof. This is now just a matter of piecing together the previous lemmas. Suppose there is a function

$$\rho = \frac{p + it}{q + is} \in R_n^m(\mathbb{C}) \quad (5.9)$$

for which

$$\|f - \text{Re } \rho\| < \|e\|. \quad (5.10)$$

We may also assume that $Z(q^2 + s^2) \cap Z(q_0) = \emptyset$. From Lemma 3.4 choose ω , α , β , δ , and γ so that

$$\omega(p_0\beta - q_0\alpha)q_0 - p_0\gamma^2 + q_0\delta = q_0[pq - st] - p_0[q^2 + s^2], \quad (5.11)$$

and construct r , r_λ , L , and L_λ as in equations (4.1)–(4.4). By Lemma 2.3 we have that for a sufficiently large c

$$\|e_c - \text{Re } L\| < \|e_c\|. \quad (5.12)$$

From Lemma 2.2 we can in fact assume

$$\|e_c - L\| < \|e_c\|. \quad (5.13)$$

Replacing f by f_c for some large c , we may assume that

$$\|e - L\| < \|e\|. \quad (5.14)$$

From Lemma 5.3, r_0 is not a local best approximation to f . ■

ACKNOWLEDGMENT

I am grateful to Professor Li Wu from Hangzhou University for several insightful comments, which in particular helped formulate Section III.

REFERENCES

1. E. W. CHENEY, "Introduction to Approximation Theory," McGraw-Hill, New York, 1966.
2. M. H. GUTKNECHT, Non-strong uniqueness in real and complex Chebyshev approximation, *J. Approx. Theory* **23** (1978), 204–214.
3. M. H. GUTKNECHT AND L. N. TREFETHEN, Real and complex Chebyshev approximation on the unit disk and interval, *Bull. Amer. Math. Soc.* **8** (1983), 455–459.
4. G. MEINARDUS, "Approximation of Functions Theory and Numerical Methods" (translated by L. Schumaker), Springer-Verlag, New York, 1967.
5. D. J. NEWMAN, "Approximation with Rational Functions," Regional Conference Series in Mathematics, Amer. Math. Soc., Providence, RI, 1979.
6. E. B. SAFF AND R. S. VARGA, Nonuniqueness of best approximating complex rational functions, *Bull. Amer. Math. Soc.* **83** (1977), 375–378.
7. D. WULBERT, The rational approximation of real functions, *Amer. J. Math.* **100** (1978), 1281–1315.
8. D. WULBERT, The characterization of complex rational approximations, *Illinois J. Math.* **24** (1980), 140–155.